

MNCH Indicators Pre/Post MamaToto Intervention - Bushenyi District

Indicator	2012 Pre	2014 Post	Absolute Change (%)
Health Practices			
Improved Water Source	82%	91%	9%*
Improved Latrine/Sanitation	59%	71%	12%
U5 Bednet Use	70%	79%	10%*
Vitamin A supplementation in last 6 months (6-23 months)	63%	76%	12%*
Dewormed in last six months (12-59 months)	62%	82%	20%*
DPT3 (12-23 months)	83%	86%	4%
Measles (12-23 months)	74%	83%	10%*
Exclusive Breastfeeding (0-5 months)	49%	60%	11%
Care Seeking/Health Access			
Fever; antimalarial treatment	34%	35%	1%
Diarrhea; ORS treatment	37%	33%	-4%
Presumed pneumonia; antibiotics given	50%	67%	17%*
Four or More ANC Visits	60%	73%	13%*
Skilled attendant present at delivery	78%	88%	10%*
Delivery at Health Facility	77%	85%	8%*
Postnatal Visit	43%	80%	37%*
Postnatal Visit <48 hours	32%	75%	43%*
Contraceptive Prevalence Rate	55%	63%	9%*
Unmet need for contraception	40%	24%	-15%
VHT Encounters			
Home visit past year	24%	65%	41%*
Advice for sick child	22%	47%	24%*
Health education session past year	24%	31%	7%*
Morbidity			
Fever	23%	20%	-3%
Diarrhea	19%	13%	-7%*
Presumed Pneumonia	22%	18%	-4%*
U5 Underweight (Moderate & Severe)	17%	14%	-3%*
U5 Stunting (Moderate & Severe)	36%	35%	-1%
U5 Wasting (Moderate & Severe)	8%	6%	-2%

*statistically significant

[Source: Baseline (2012) and Endline (2014) Study]