

MNCH Indicators Pre/Post MamaToto Intervention - Bushenyi District

Indicator	2012 (%) Pre	2014 (%) Post	Absolute Change (% (95% CI))
Health Practices			
Improved Water Source	82.5	91.1	8.6 (4.4, 12.7)
Improved Latrine/Sanitation	59.4	71.1	11.7 (-0.2, 23.6)
U5 Bednet Use	69.6	79.2	9.6 (3.7, 15.5)
Vitamin A supplementation in last 6 months (6-23 months)	63.1	75.6	12.5 (4.0, 21.0)
Dewormed in last six months (12-59 months)	62.0	82.4	20.4 (12.2, 28.5)
DPT3 (12-23 months)	82.6	86.2	3.6 (-4.9, 12.1)
Measles (12-23 months)	73.6	83.2	9.6 (1.3, 17.9)
Exclusive Breastfeeding (0-5 months)	49.1	59.8	10.7 (-3.8, 25.2)
Care Seeking/Health Access			
Fever; antimalarial treatment	34.4	35.0	0.6 (-9.4, 10.6)
Diarrhea; ORS treatment	37.5	33.1	-4.4 (-15.2, 6.4)
Presumed pneumonia; antibiotics given	50.0	67.2	17.2 (7.7, 26.7)
Four or More ANC Visits	59.7	72.7	13.0 (7.5, 18.4)
Skilled attendant present at delivery	78.3	88.4	10.1 (5.6, 14.5)
Delivery at Health Facility	77.3	85.3	8.0 (2.7, 13.3)
Postnatal Visit	42.6	80.0	37.4 (31.3, 43.4)
Postnatal Visit <48 hours	32.5	75.1	42.6 (38.1, 47.0)
Contraceptive Prevalence Rate	54.6	63.3	8.7 (4.4, 12.9)
Unmet need for contraception	39.6	24.4	-15.2
VHT Encounters			
Home visit past year	23.9	64.6	40.7 (32.5, 48.9)
Advice for sick child	22.3	46.6	24.3 (15.3, 33.3)
Health education session past year	23.9	31.2	7.3 (1.7, 12.9)
Morbidity			
Fever	23.1	20.0	-3.1 (-8.1, 1.9)
Diarrhea	19.2	12.6	-6.6 (-10.1, -3.1)
Presumed Pneumonia	22.3	17.9	-4.4 (-8.5, -0.3)
U5 Underweight (Moderate & Severe)	16.9	14.0	-2.9 (-6.4, -0.6)
U5 Stunting (Moderate & Severe)	36.3	35.4	-0.9 (-7.2, 5.4)
U5 Wasting (Moderate & Severe)	7.9	6.4	-1.5 (-3.8, 0.8)

[Source: Baseline (2012) and Endline (2014) Study]